

Biological Impacts of Climate Change

John P McCarty, *University of Nebraska at Omaha, Omaha, Nebraska, USA*

L LaReesa Wolfenbarger, *University of Nebraska at Omaha, Omaha, Nebraska, USA*

James A Wilson, *University of Nebraska at Omaha, Omaha, Nebraska, USA*

Based in part on the previous version of this eLS article 'Biological Impacts of Climate Change' (2009) by John P McCarty, L LaReesa Wolfenbarger, James A Wilson.

Introductory article

Article Contents

- Introduction
- How Climate Impacts Life
- Observations of the Biological Impacts of Climate Change
- Trying to Understand the Future
- Summary

Online posting date: 20th March 2017

Climate has far reaching impacts on biological systems. Survival and reproduction depend on how well adapted individuals are to local climate patterns. Climate change can disrupt the match between organisms and their local environment, reducing survival and reproduction and causing subsequent impacts on populations or species' distributions across geographic regions. Changes in climate may benefit some species and cause extinction for others. Cumulatively, it will alter biological communities and the functioning of ecosystems. Changes to ecosystem functions can in turn increase or decrease the rate of human-driven climate change. In addition to effects of climate variables such as temperature and precipitation, plants may respond directly to rising concentrations of CO₂, while aquatic species cope with changes in water chemistry as greenhouse gasses dissolve in water. The earth is already experiencing sufficient climate change to affect biological systems; well-documented changes in plant and animal populations are related to recent climate change. Predicting future biological impacts of climate change remains a formidable challenge for science.

Introduction

Emission of carbon dioxide, methane and other greenhouse gasses is a primary driver of current and future human-caused climate change (IPCC, 2014). Biological systems respond not only to the resulting changes in climate but also directly to

changes in atmospheric concentrations of greenhouse gasses and to secondary effects such as rising sea levels and changes in water chemistry.

One of the fundamental lessons from the science of ecology is that patterns of climate strongly influence the distribution and abundances of living organisms. Climate describes weather patterns for a given location over an extended period of time (e.g. 10+ years). Climate is not just the average conditions of temperature and precipitation but also the seasonal and annual weather variation, including the frequency and severity of extreme events such as storms or drought. The earth's climatic patterns are now changing and human activities are major contributing factors driving these changes (IPCC, 2014). The types of impacts climate change will have on living systems have far reaching consequences for natural ecosystems and the people who depend on the goods and services that ecosystems provide.

The distribution of the general types of plants and animals, or biomes, around the world can be predicted with some accuracy from climate patterns. General climate patterns are in turn generated by atmospheric circulation patterns caused by differential heating of the earth's surface, which determine the temperature and precipitation patterns in a given area. Relatively minor changes in the overall heat balance of the earth can change atmospheric circulation and result in local climate changing more dramatically than indicated by the degree of average warming. For example, a 1°C increase in average temperature can translate into significantly longer growing seasons, greater number of extremely hot days, and changes in the patterns and intensity of precipitation (Melillo *et al.*, 2014). In addition, a 1°C increase in global average temperature will result in some geographic regions experiencing much greater than 1°C warming, while other regions show little or no warming. This has been observed with the warming of recent decades where the arctic has warmed much more quickly than the global average (IPCC, 2014). **See also: [Ecosystem Concepts: Introduction](#); [Plant Physiological Responses to Climate and Environmental Change](#)**

Climate changes of this magnitude have already been observed to impact species and entire biological communities (McCarty, 2001, Rosenzweig *et al.*, 2008). Ongoing and future climate changes will have far reaching effects on biological systems, including humans. Each species is likely to respond differently to changes in climate. For some species, climate will remain within

eLS subject area: Ecology

How to cite:

McCarty, John P; Wolfenbarger, L LaReesa; and Wilson, James A (March 2017) Biological Impacts of Climate Change. In: eLS. John Wiley & Sons, Ltd: Chichester.
DOI: 10.1002/9780470015902.a0020480.pub2




Figure 1 The current geographic range of species is limited to areas with a suitable climate (a). As climate changes and the areas with suitable climate shift towards the poles, species will respond in different ways. For some species, geographic distributions might shift to track changes in suitable climate, with little change in the overall size of their range (b). However, for other species the area of suitable habitat may decline (c), or their ability to shift their geographic range to take advantage of new areas may be limited by physical barriers such as mountains or bodies of water (d) or restrictions on the movement of individuals that limit the ability to disperse (e). However, other factors besides climate can influence future geographic distributions. For example, some species will evolve to adapt to new climatic conditions and remain in their current geographic range, while interactions with competitors, predators and pathogens might prevent species from using areas with newly suitable habitat (f). Finally, species that are unable to respond adequately to new climatic conditions or whose suitable habitat becomes too small (c) will go extinct. Reproduced with permission from Lambers (2015) © The American Association for the Advancement of Science.

the existing range of tolerances. If a local climate shifts outside the species' range of tolerances, one of three responses will occur: adaptation (environmental or genetic change), relocation or extinction (**Figure 1**).

One of the great challenges for biology today is to try to understand how future changes in climate will impact biological systems. Progress towards this goal depends on understanding the underlying mechanisms as to how individuals and species respond to changes in climate, examining biological responses to recent climate changes and integrating this information in experiments and models to try to understand how complex biological systems will interact under future changing climate conditions.

How Climate Impacts Life

The basic components of climate – temperature and moisture – have pervasive impacts on organisms. Physiological processes define life, and at the most basic level these are chemical reactions. As such, they are subject to the unavoidable relationship between temperature and the speed of chemical reactions. While the rates of simple chemical reactions increase as temperature increases, physiological processes respond in a more complex manner. This typically involves a thermal optima where the reaction proceeds most quickly. Physiological processes proceed more slowly at temperatures above or below the thermal

optimum. Most physiological processes are also water-based. All organisms face a major challenge: maintaining an appropriate water balance and temperature range for life-sustaining physiological processes while living in environments that are too wet/dry and too cold/warm.

The climate where an organism lives dictates the specifics of this challenge, and organisms have evolved numerous adaptations to cope with hostile environments. Organisms are exposed to not just the average temperature and moisture conditions but also the variability associated with seasons and with extreme events. Plants and animals cope with variation in the environment in many ways. Some animals move to different geographic areas to avoid severe conditions (e.g. migration). Both plants and animals can reduce activity when conditions are too severe: torpor in animals and senescence in plants. **See also: Fundamentals of Water Relations and Thermoregulation in Animals; Ecology of Water Relations in Plants**

Climate and individuals

The physiologic health or condition of individuals acts as the link between the habitat and population dynamics. Individuals poorly suited for their habitat may not obtain enough energy to maintain themselves in good condition and may forego reproduction completely or until conditions in the habitat improve.

Climate can impact survival and reproduction for animals that are able to regulate their body temperature within a narrow range even as the surrounding air temperature varies, including endotherms such as birds and mammals. As the temperature of the environment decreases relative to body temperature, individuals need to expend energy to stay warm and may eventually reach the point where they can no longer generate enough heat to maintain body temperature. If individuals cannot move to a warmer environment or obtain needed energy, they may die. At the other extreme, as environmental temperature rises above body temperature, individuals need to expend energy and often significant amounts of water, to stay cool. Once again if surrounding temperatures rise too high, the individual will no longer be able to regulate the body temperature and could die. Even if environmental temperatures do not reach the extremes that cause death, the energy required to maintain internal conditions is no longer available for other important activities such as reproduction.

Climate also has profound effects on the survival and reproduction of ectotherms: animals such as amphibians, reptiles and insects that are unable to regulate their body temperature independently of the surrounding temperature of the environment. For these species, the rate of physiological processes determined by body temperature depends upon ambient temperature. As ambient temperature rises the rate of physiological processes increases in a nonlinear fashion and increases more rapidly when initial temperatures are low. For example, a rate may triple across a temperature of 10–20°C but may only double from 20 to 30°C. Although ectotherms have a limited ability to avoid the physiological consequences of changing environmental temperature, they have evolved a broad assortment of strategies for coping with a broad range of climate conditions. **See also: Thermoregulation in Vertebrates; Vertebrate Metabolic Variation**

As an adaptation to harsh environmental conditions, especially cold temperatures, some ectotherms become inactive during unfavourable seasons of years, and climate change will significantly alter the energy expenditures and body condition of these organisms. Body condition in ectotherms is tightly bound to reproductive output, timing to maturity and survival during inactive periods. Unlike hibernation in mammals, where individuals can regulate body temperature independent of ambient temperature, ectotherms cannot. Because their hibernating metabolic rate is dependent on ambient temperature, warmer winters will cause ectotherms to utilise more energy during hibernation than colder winters. For example, common toads (*Bufo bufo*) in the United Kingdom have shown a decline in body condition associated with increasing temperatures which is also associated with decreased annual survival (Reading, 2007). In addition, when ectotherms use more energy to hibernate during warming winter, they emerge from hibernation in poorer condition, which reduces reproduction. **See also: Hibernation: Endotherms; Hibernation: Poikilotherms; Thermoregulation in Vertebrates: Acclimation, Acclimatization and Adaptation**

Aquatic species are also sensitive to temperature. Distributions of many species are limited by water temperatures. The tolerance limits for some species can be quite narrow. Corals, for example, can lose their symbiotic algal partners with warming on the order of 1°C, resulting in bleaching that stresses or even kills coral (Doney *et al.*, 2012). In addition, the dissolved

oxygen levels decline as water temperature rises, placing an additional limit on some species. Antarctic icefish (Notothenioidei: Channichthyidae) provide an extreme example of how trade-offs between temperature and dissolved oxygen availability impact aquatic organisms. This group of fish has lost functioning haemoglobin due to the loss of the gene that produces the β -subunit of the haemoglobin protein (Cocca *et al.*, 1995). This mutation would normally prove to be lethal, resulting in the metabolic asphyxiation of the individual because nonfunctional haemoglobin cannot carry oxygen to the body's tissues to support aerobic metabolism. However, the fact that they live in extremely cold Antarctic waters results in low metabolic rates and a substantial reduction in oxygen demand. Indeed, the waters are so cold that icefish depend on several physiological adaptations to help them survive, including antifreeze proteins in their blood. Icefish live within a few degrees of their upper, lethal temperature limit and if these fish were ever exposed to warmer waters their metabolism, and consequently their oxygen demand, would increase resulting in their death (Bilyk and DeVries, 2011; Beers and Sidell, 2011). Currently, the waters surrounding Antarctica have increased in temperature from 0.17°C to ~1°C over the last 50 years and are predicted to increase by another 2°C throughout the next century (Clarke *et al.*, 2007).

Plant physiology is also sensitive to temperature range and moisture balance. Temperature and moisture interact to determine the rate of photosynthesis, the physiological process in which plants meet their energy needs and use the sun's energy to synthesise carbohydrates from carbon dioxide and water. Because plants have limited ability to regulate internal temperatures or to avoid temperature extremes, many species become inactive during seasons when conditions are unfavourable. Therefore, climate determines the length of the growing season and the nongrowing season when plants lack the necessary moisture for photosynthesis or when temperatures drop below the freezing point of water.

The response of plants to global change is complicated by the fact that one of the primary greenhouse gasses, carbon dioxide, is also the key building block in photosynthesis. Rising concentrations of carbon dioxide will increase rates of plant productivity, but only when sufficient water and nutrients are available. The net impact of global change on photosynthesis will vary regionally, with water limitations and extreme temperatures counteracting the positive impacts of higher carbon dioxide concentrations in many areas. **See also: Ecology of Water Relations in Plants; Photosynthesis: Ecology; Plant Physiological Responses to Climate and Environmental Change; Plant Responses to Elevated CO₂**

Climate and populations

A population consists of a collection of individuals, and population size changes due to reproduction, immigration, mortality and emigration of these individuals. As such, a population can only grow when net individual reproduction and immigration is greater than mortality and emigration, and decreases when the opposite occurs. Population size is influenced by a complex interaction of direct and indirect factors that change the energy budget of individuals living in a population. Simply put, direct factors are those that are abiotic and reflect changes to a population due to thermal

stress, extreme weather or changes in precipitation. Indirect factors represent changes to the biotic environment, typically manifested as changes in biotic interactions due to resource availability, community composition and structure and predation pressure.

Most species have resolved the challenges associated with a specific range of climate conditions, and the occurrence of these conditions constrains the geographic range of a species. Outside of this range, factors driving populations down – mortality and emigration – will overwhelm reproduction and immigration, and populations will die out. In some cases, the correlation between specific climate conditions and the limits of geographic range can be quite close. Classic examples where climate limits geographic range include plants such as wild madder (*Rubia perigrina*) whose northern limit in Europe corresponds closely to where January temperatures remain approximately 4.5°C and saguaro cactus (*Carnegiea gigantea*) in Arizona where the range is limited to areas where daily temperatures consistently rise above freezing (Begon *et al.*, 2006). The distribution of most species is influenced by a more complex combination of factors, though climate may still play a role. For example, the distribution of Northern Bobwhite (*Colinus virginianus*) in North America is explained in part by a combination of temperature and precipitation (Matthews *et al.*, 2007; **Figure 2**).

Although unfavourable climate can eliminate populations, favourable climate is not a guarantee that populations will occur or persist. Additional environmental conditions besides climate may be unfavourable, or geographic barriers may exist that prevent members of the population from ever reaching a given geographic region. **See also: Range Limits**

Climate and interactions among species

Individuals interact with members of other species in a variety of ways. These include interactions among parasites, disease organisms, predators and prey and competition over common resources as well as mutually beneficial interactions such as flowering plants producing nectar for the animals that pollinate their flowers. Through these interactions species may influence the population size and even the geographic range of other species. Interactions among species in a community can produce indirect links between a population and climate. For example, rush moths (*Coleophora alticolella*) in England tolerate the direct effects of climate associated with high elevations sites, but the plants on which the caterpillars feed do not produce enough seeds at high elevation, thereby limiting the range of the moth to low elevation sites with abundant food for caterpillars (Randall, 1982). Differences among species in their abilities to cope with changes in climate can shift the balance between competitors or allow new predators, pathogens or parasites to invade a species range. **See also: Coexistence; Community Ecology: An Introduction; Interspecific Competition; Interspecific Interaction; Predation (Including Parasites and Disease) and Herbivory**

Climate and communities and ecosystems

The community of plants and animals in a given area emerges from the responses of individual species to climate and other

physical factors. The species-specific nature of the factors limiting ranges means that biological communities do not respond to climate as cohesive units but rather are assemblages that reflect the tolerances of their component species. These assemblages of species are both familiar and have emergent properties such as diversity and productivity. As climate changes it is probable that assemblages of species that are now familiar will be broken up as species that respond to new environmental conditions in different ways. New assemblages will then emerge with their own characteristics and properties. **See also: Community Ecology: An Introduction**

Communities in turn are linked with the nonliving environment in ecosystems. Within an ecosystem, the flow of energy and matter among organisms is constrained by the ability of plants to capture the sun's energy in a form that can be used by other organisms. In this way, energy balance forms the link between individual condition, population dynamics and ecosystem functioning. Likewise, these higher-level processes that emerge from ecosystems have far reaching implications for humans through their impacts on nutrient, air and water cycles. **See also: Ecosystem Concepts: Introduction; Photosynthesis: Ecology**

Observations of the Biological Impacts of Climate Change

The earth's climate has been in a state of change for most of the history of life, and ample evidence exists to show how biological systems respond to changes in temperature and moisture. We know from fossils and other remains of long dead organisms that biological systems have undergone dramatic changes in response to past changes in climate. **See also: Palaeoclimatology; Palaeoenvironments**

Since the mid-twentieth century, climate has been changing rapidly relative to previous periods. Global mean temperatures have increased by about 0.85°C since 1880 and this warming has occurred unevenly across the globe (IPCC, 2014). Temperatures over land are warming faster than over oceans, and warming of the arctic is occurring almost twice as fast as the global average.

Other changes in the environment follow from this warming. Polar regions have had extensive warming of air and surface ocean temperatures. This has led to a decrease in the extent of Arctic sea ice, especially during summer. Similarly, ice mass has been lost from the Greenland ice sheet and the northern Antarctic Peninsula (IPCC, 2014). In the Northern Hemisphere, the extent of snow cover in the spring months of March and April has declined at a rate of 1.6% per decade since the mid-twentieth century (IPCC, 2014). These changes both reflect a changing climate and can impact species that are dependent on ice and snow or whose life cycles are timed to a predictable spring thaw.

The magnitude of recent changes are sufficient for scientists to observe how biological systems respond to ongoing climate change. Researchers have turned to long-term datasets to understand how climate change is impacting biological systems. Some of the longest running quantitative records of biological systems are observations of seasonal biological events. In many areas of the globe, records of the timing, or phenology, of events such




Figure 2 Much of the geographic range of species can be explained by climate. In the example shown above, Northern Bobwhites are widespread across the southern two-thirds of the eastern United States. (a) The map shows their geographic range and relative abundance based on the North American Breeding Bird Survey. USDA Forest Service scientists evaluated the geographic distribution of Northern Bobwhite (and 146 other bird species) against information about the climate and vegetation in the eastern United States. The importance of different climate and habitat variables in explaining geographic range depends on the bird species. (b) Illustration shows the combination of temperature and precipitation found in the eastern United States. The coloured cells indicated the combinations of temperature and precipitation where Northern Bobwhites are found, while the light grey squares represent combinations of temperature and precipitation found in the eastern United States where Northern Bobwhites are absent. Reproduced from Matthews *et al.* (2007) © USDA Forest Service.

as when birds migrate and when leaves or flowers appear in the spring go back decades or more. These data provide strong evidence for widespread changes in the timing of biological events linked to the climate changes experienced over the same time period. For example, Terry Root and her colleagues conducted a review of 64 studies of biological events associated with the onset of spring in 694 species of plants and animals; the results indicated that spring phenological events like migration and flowering have been occurring earlier at a rate of 2–7 days every decade (Root and Hughes, 2005). Globally, the documented changes in biological systems are consistent with the changes in climate observed during these same decades (Rosenzweig *et al.*, 2008). **See also: Climate Change Impacts: Insects; Climate Change Impacts: Birds**

Not all species will be able to cope with climate change by tracking favourable conditions because the changing climate might exceed the capacity of individuals to alter the timing of activity. Desert-dwelling ground squirrels face extreme heat during summer months and also low water availability. As a result, ground squirrels enter a period of seasonal inactivity (aestivation) during the summer months to decrease energetic demands. However, as summers become warmer and precipitation decreases, the length of the inactive season will increase while the wet season, when squirrels obtain food for aestivation, shortens. Piute ground squirrels (*Spermophilus mollis*), from the western United States, have been shown to enter aestivation early in response to early spring drought. If the drying trend continues to increase, these animals may not acquire sufficient energy resources during the shortened growing seasons to aestivate over the longer dry months (Steenhof *et al.*, 2006).

Changing phenology may not always track favourable conditions if it disrupts or decouples species interactions or other biotic factors that influence populations. For example, Marmots in the mountains of Colorado spend the winter hibernating and emerge from hibernation when air temperature warms. As air temperatures have increased, marmots have been emerging from hibernation 23 days earlier than they did in 1976. However, because winter precipitation in the form of snow is increasing, deep snow is still on the ground preventing plants from growing. Therefore, marmots must use their limited fat reserves for longer periods in spring while at the same time they have increased energetic demands and are preparing for reproduction (Inouye *et al.*, 2000).

Geographical ranges of species have expanded and contracted as climate factors have changed over the past decades. Historic datasets show that geographic ranges of many northern hemisphere species are shifting northward at rates consistent with observed climate change. A recent meta-analysis examined the results from studies of looking at changes in latitudinal distribution of taxonomic groups covering 764 species in Europe, North America and Chile and found that the groups had shifted away from the equator at a median rate of 16.9 km per decade (Chen *et al.*, 2011). Overall, the distance moved corresponded to the expected change given the rate of climate warming. However, the distance moved varied among species groups and some groups were not moving fast enough to keep up with the pace of climate change. In mountainous regions an analogous shift has been documented with species spreading to higher elevations where temperatures are cooler at a median rate of 11.0 m per

decade (Chen *et al.*, 2011). These range expansions lead to a homogenisation of alpine plant communities and a decrease in specialised communities, particularly in alpine tundra associated with mountain peaks (Jurasinski and Kreyling, 2007). Marine species are also altering their ranges to compensate for increased water temperatures. Globally, marine species have shifted their distribution 30.6 ± 5.2 km per decade (Sydesman *et al.*, 2015). Most range shifts seem to be associated with an attempt to maintain their presence in cooler waters and have therefore resulted in more polar shifts, or, when polar movements are not possible (i.e. movement is blocked by the presence of land), then species may move to deeper waters. **See also: Alpine Ecosystems; Marine Communities**

Most species are likely to respond to changing conditions in more than one way. For example, Pacific walrus use sea ice as a place to rest between foraging trips, a location to molt their skin, and as a safe location to give birth to their young (Jay *et al.*, 2012). In response to earlier ice break up in spring and later ice formation in fall, walrus now begin their spring migration about 1 month earlier and return in the fall a month later (MacCracken, 2012). In addition, populations in the Chukchi Sea have shifted their range to more northern latitudes in June and July, while during September and October walrus forage closer to shore than historically documented. Even with these adjustments, the loss of sea ice has contributed to young and adult females to haul out in large numbers onto the shore (Jay *et al.*, 2012; MacCracken, 2012).

In some cases it appears that changes in climate have overwhelmed species' abilities to respond. Amphibians living in the mountains of Central America may provide a window into what the future might hold (Pounds *et al.*, 2006). The Monteverde region of Costa Rica is world famous for its cloud forest and diverse community of frogs. Declines in populations of these frogs have been well documented in recent decades and include the extinction of the golden toad (*Bufo periglenes*). Population decline and species loss are linked to the changing climate, including a reduction in misty days; a defining feature of cloud forests, but the full explanation is much more complex. Warmer night time temperatures produce ideal growing conditions for a chytrid fungus (*Batrachochytrium dendrobatidis*) that infects amphibians and sometimes kills them. Frogs stressed by warmer and drier climate are less able to resist fungal infection. As a result, populations decline and populations and species go extinct (Pounds *et al.*, 2006).

Responses of a few key species to climate change may result in far reaching consequences for entire ecological communities. Corals are sensitive to ocean warming, which contributes to bleaching. In addition, increases in atmospheric CO₂ concentrations result in more acidic ocean waters as CO₂ dissolves in water. These more acidic conditions make it more difficult for corals to secrete their skeletons (Doney *et al.*, 2012). Loss of corals and the physical structures they create will result in disruption of the entire community dependent on this habitat. These changes are already impacting many of the world's reef systems and global impacts are projected in the coming decades (Frieler *et al.*, 2013). An analogous change has been observed along the temperate coast of Australia where extreme warming starting in 2011 resulted in the loss of kelp forests and the community that depends on the kelp (Wernberg *et al.*, 2016).

Within 2 years, the kelp forest community had been replaced by tropical and subtropical species along 100 kms of the coast.

In addition to changes in populations and biological communities, earlier spring warming and changes in summer climate have produced significant changes in the fundamental energy dynamics of ecosystems. Satellite and ground-based observations show that over the past 30 years the thermal potential growing season in northern ecosystems has become 10.5 days longer; however, limited moisture and light has limited the ability of plants to take advantage of the warmer conditions so the growing season when photosynthesis occurs has lengthened by 6.7 days (Barichivich *et al.*, 2013). For some ecosystems this has been accompanied by an increase in net primary productivity as well. Globally, satellite data measuring leaf-area index shows that an overall 'greening' of the earth's surface has occurred during the growing season, indicating that plants produce more leaves (Zhu *et al.*, 2016). Models indicate that this increase is driven by the fertilising effect of CO₂ combined with changes to climate, changes to nitrogen cycling and other variables. Analogous changes are seen in aquatic systems (Burrows *et al.*, 2011) where spring events such as migrations and plankton blooms have occurred 4.4 ± 1.1 days earlier (Poloczanska *et al.*, 2013).

Trying to Understand the Future

Biological responses to future climate change will depend on the rate and magnitude of continued climate change. Sophisticated models of the earth's climate system incorporate our best understanding of the variables that interact to drive the earth's climate, but a fundamental uncertainty for predicting climate change is how human behaviour will influence the increase in atmospheric greenhouse gas concentrations. The Intergovernmental Panel on Climate Change (IPCC) presents the most comprehensive and authoritative synthesis of possible outcomes by examining several leading climate models and incorporating a range of 'scenarios' for future greenhouse gas concentrations. The IPCC's 2014 report suggests that warming will continue and accelerate during the twenty-first century. The average global temperature by the end of this century is projected to increase 0.3–4.8°C or more beyond what was experienced during 1986–2005 (or approximately 0.9–5.4°C above pre-industrial levels), depending on how human emissions of greenhouse gases change (IPCC, 2014). As with past change, actual changes in temperature experienced by plants and animals will vary. Alpine and polar regions have experienced the greatest change in climate and will likely continue to show dramatic effects of global warming. **See also: Biotic Response to Climatic Change**

Methods for studying the biological impacts of climate change – models and experiments

One approach to understanding complex interactions among species and a changing environment is to take what we know about how species respond to climate (Figure 2) and combine that information with output from global climate models to forecast

how species or communities might respond to future conditions. For example, the USDA Forest Service has detailed information about where different forest types currently exist (Prasad *et al.*, 2007; Figure 3). By analysing these distributions in relation to environmental variables, predictive models can be generated that approximate the distribution currently observed ('current modelled' in Figure 3). By then, incorporating forecasts of how those environmental variables will change under scenarios of incorporating with low or high rates of climate change, the future distributions of the forest types can be generated. In the case shown in Figure 3, spruce/fir forests currently exist in northern United States but the models suggest that future conditions will no longer be suitable for that forest community. Although these models do not predict whether species will be able to respond to climate change successfully, they can provide insights into how potential habitat for different species might shift under different climate conditions.

Another approach to understanding complex interactions is to expose current communities to conditions they might expect in the future. For example, experiments can alter temperature, moisture levels, concentrations of carbon dioxide in the surrounding air and various combinations of these and other variables (Figure 4). This approach can be especially useful for understanding how changes in different aspects of the environment can interact to influence biological systems. For example, the Aspen FACE (Free-Air Carbon Dioxide Enrichment) experiment exposed trembling aspen (*Populus tremuloides*) and other trees to higher concentrations of carbon dioxide as well as the air pollutant ozone to understand impacts on forest productivity and ecosystem function (Karnosky *et al.*, 2003). The TasFACE project employed similar technology to expose native grasslands in Tasmania, Australia to increased levels of CO₂ and then added treatments including infra-red heaters to mimic higher temperatures expected in 2050, (Hovenden *et al.*, 2014; Figure 4). **See also: Global Carbon Cycle; Climate Change Impacts: Vegetation**

Effects of increasing carbon dioxide on biological systems

Most of our attention is focused on changes in climate driven by increasing concentrations of carbon dioxide in the atmosphere. However, carbon dioxide can have other effects on biological systems that will contribute to changes in the coming decades.

Carbon dioxide plays a key role in photosynthesis. Photosynthesis rates can be higher as carbon dioxide concentrations increase. Whether plants are able to take advantage of the enrichment of the atmosphere with carbon dioxide will depend on whether they have the other resources, such as water and nutrients, to support increased productivity. The TasFACE experiments (Figure 4) demonstrated that variation in seasonal rainfall was a factor that determined how grasslands respond to elevated CO₂, in part because of interactions between rainfall in the cool months and limited nitrogen (Hovenden *et al.*, 2014). Other anthropogenic factors can also influence these patterns. For example, the aspen trees in the Aspen FACE project that were grown under higher concentrations of carbon dioxide grow faster, whereas those exposed to increased ozone grow slower (Karnosky *et al.*, 2003; Figure 4). When exposed to both carbon




Figure 3 Mathematical models provide one approach for helping us understand how changes in climate will impact biological systems. These maps show the current geographic range of forest types as well as modelled output based on current climate and two scenarios of future climate. Current forest types (panel a) are based on the USDA Forest Service's Forest Inventory Analysis (FIA) data. Information about the geographic range of 134 tree species was evaluated against 38 environmental variables to generate predictive models. The utility of the models can be evaluated by inserting current climate conditions into the models and comparing the output (panel b) to current distributions of forest types (panel a). The general correlation between the actual current FIA data and the modelled current distributions indicates that much of the variation where the forest types occur can be explained by combinations of climate variables. This correlation also suggests that the Forest Service model can be used to model potential habitats under future climate conditions. The scientists took the output from three widely used global climate models under two scenarios used by the Intergovernmental Panel on Climate Change. The 'Low' scenario assumes that emissions of greenhouse gases will be significantly reduced, while the 'High' scenario assumes that current emission trends will continue. Panels (c) and (d) show how the potential habitat for forests might change in the future. Note in particular the loss of potential habitat for northern forest types such as Spruce-Fir forests that are currently found in the northern tier of states but which might disappear in the future. Reproduced from Prasad *et al.* (2007) © USDA Forest Service

dioxide and ozone the effects on growth were neutralised. Indirect effects on trees appeared when trees were exposed to higher levels of carbon dioxide by impacting insects that live and feed on the trees and by altering the competitive balance between aspen and sugar maple trees (*Acer saccharum*). Not all plants in a community will be equally able to respond to increased carbon dioxide. Plants that are able to respond may have a competitive advantage

in communities. In the grasslands of North America, this might lead to invasion of grasslands by trees and shrubs as woody plants gain an advantage over grasses. Exposure to higher levels of CO₂ during growth can also change the nutrient levels of plants in ways that could have implications for the animals that eat them. In Australia, the leaves of rainforest trees grown under elevated CO₂ contained lower levels of nitrogen and other nutrients (Kanowski,




Figure 4 Experimental approaches for studying the effects of climate change on biological systems. (a) Species may respond directly to warming associated with climate change. Open Top Chambers are designed to passively warm vegetation plots with a simple, inexpensive system that can be replicated across many sites as part of the International Tundra Experiment (ITEX; Elmendorf *et al.*, 2012). Reproduced with permission from R Hollister. (b) Changes in precipitation patterns can be manipulated using shelters such as these deployed in a salt marsh as part of study to study the response of ecosystem processes such as plant growth and nutrient cycling. Reproduced with permission from H Emery. (c) Plants may respond directly to changes in concentrations of greenhouse gasses such as carbon dioxide. The Aspen FACE (Free Air Carbon Dioxide Enrichment) Experiment exposed trembling aspen trees in the open under carbon dioxide levels similar to those expected to occur late in the twenty-first century. The pipes surrounding the growing trees release carbon dioxide, mimicking the effects of altered atmosphere in a field setting where plants interact with each other and with other environmental variables in a natural setting. Photo by JP McCarty. (d) The TasFACE experimental system combines the FACE technology with infrared heaters to simulate warming and altered atmospheric gasses simultaneously. Reproduced with permission from M Hovenden.

2001). A decline in the nutritional value of the leaves in turn raises concerns about the impacts on the native marsupials that feed on the leaves. **See also: Photosynthesis: Ecology; Plant Physiological Responses to Climate and Environmental Change; Ecophysiological Responses of Plants to Air Pollution**

Increased carbon dioxide in the atmosphere is also having far reaching direct effects on the world's oceans. Carbon dioxide is soluble in water and decreases its pH by forming carbonic acid. Oceans have become more acidic in recent years as atmospheric carbon dioxide levels have increased. The average pH of the ocean has decreased from a pre-industrial level of 8.20 to 8.11 and could fall to 7.8 by 2100 (Feely *et al.*, 2009). Many aquatic organisms are sensitive to the acidity of water. For example, the carbonate shells of marine animals can dissolve in acidic

water. Increased acidity will further stress coral reef communities that are already suffering the effects of warmer water temperatures. **See also: Climate Change and Biogeochemical Impacts; Global Carbon Cycle**

Range shifts in response to climate change

As climate changes, the geographic range of some species will shift to track changes in climatic conditions (**Figure 1**). The total area occupied by a species might increase, decrease or remain constant. The potential for the range of species to move with the climate will depend on whether there is a net gain or loss of area with a suitable climate. Some species whose ranges are currently

limited by climate may expand into suitable areas whereas the potential range of other species will shrink. At a local scale, suitable climate conditions may disappear entirely. This might be most dramatic in mountains where the potential shift in habitats to higher elevations is limited by the height of the mountains or near the poles where sea ice habitat will disappear. Within a given area some species will be lost as the climate becomes unsuitable but other species will be added as climate becomes favourable for them. Across North America, changes in climate will result in both gains and losses of birds (Langham *et al.*, 2015; **Figure 5**). In addition, the species within a community are likely to respond differently so the composition of communities will change.

Another key factor will be the balance between how fast climate shifts and how quickly species can respond. How far individuals or seeds and other propagules can spread or move will limit changes in geographic range or migration (**Figure 1**). Freshwater plants and animals may have even more limited abilities to colonise new habitats as lakes and streams warm. Although we might predict distributions of highly mobile species to change quickly, a suitable habitat may depend not just on climate conditions but also on the presence or absence of other species or resources. Barriers to movement also exist in the form of mountains and water bodies (or land in the case of aquatic species). Finally, much of the earth's surface has been transformed by human activities, and the ability of populations of plants and animals to colonise new areas in a human-dominated landscape is uncertain. **See also: Dispersal; Biogeography; Range Limits**

Evolution in response to climate change

Changes in behaviour, timing of life history events (phenology), selective use of more hospitable microhabitats within the existing range and changes in physiology can all help species adjust to variations in climate. For some species, these capabilities already exist in individuals and responding to climate change will be relatively straightforward (i.e. phenotypic plasticity). In other cases, the characteristics that determine the climate tolerance of a species will involve a more fundamental, genetically based change (evolution).

The rate at which populations evolve or undergo change in its genetic make-up will determine whether a species can adapt to new climatic conditions. Although evolution can happen rapidly, not all species will be equally able to evolve in the face of climate change. For evolution to occur rapidly, genetic variability in traits of interest needs to exist in the population; otherwise the rate of change would depend on a suitable mutation arising, an extremely rare event. Furthermore, small or declining populations are less likely to adapt to changing environments due to their low levels of genetic variation. The rate at which evolutionary change could proceed is also a function of the time between generations. Long-lived species such as trees will evolve more slowly than species with short generation times such as insects. **See also: Adaptation and Constraint: Overview; Adaptation and Natural Selection: Overview; Natural Selection: Responses to Current (Anthropogenic) Environmental Changes; Evolutionary Responses to Climate Change**

Extinction in response to climate change

Not all species will be able to respond to changes in climate. Species that are unable to respond quickly enough will go extinct. Estimates of the risk of extinction due to climate change vary, but the magnitude of extinctions could be immense. One recent meta-analysis synthesised 131 published estimates of extinction risks and concluded that 7.9% of species were at risk of extinction although, depending on the assumptions used, some authors produced estimates as high as 50% (Urban, 2015). Extinction risks were greatest in South America, while fewer species in North America and Europe are considered to be at higher risk. Arctic species may also be at greater risk, both due to the faster warming experienced at high latitudes, and the structural changes associated with loss of sea ice.

Climate changes may increase the extinction rates of species already at risk. Small populations will have limited potential to evolve in response to new conditions. Climate change may also exacerbate the very conditions that placed species at risk in the first place. For example, invasive species, disease and parasites pose a threat to many populations, and the changing climate may facilitate the invasion of new threats into the range of species at risk. Invasive species are defined in part by their ability to colonise new areas and would be expected to expand their ranges quickly in response to climate change. This pattern has already been observed in alpine areas of Australia, where warming temperatures have allowed invasive, non-native mammals including rabbits and horses, to access areas where they had previously been excluded (Hughes, 2003). Lack of suitable habitat places many species at risk of extinction. If climate change reduces habitats further, species may be unable to recover from the loss and will go extinct. This threat is exacerbated by the fact that current conservation efforts may miss many of the species that will be at risk from climate change in the near future (Langham *et al.*, 2015). **See also: Biodiversity–Threats; Extinction**

Climate – ecology feedbacks

Climate clearly impacts biological systems but biological systems can also influence climate by changing the amount of heat absorbed from the sun, releasing water vapour and altering the levels of carbon dioxide and other greenhouse gases in the atmosphere. The impact of this feedback on climate is a considerable source of uncertainty for projecting future effects of climate change on biological systems. Plants, both on land and in the ocean, absorb carbon dioxide from the atmosphere during photosynthesis. Some of the carbon trapped this way is transferred to higher trophic levels when animals eat plants, some remains bound up in living or dead plant matter and some is released back into the atmosphere when dead plant material decays. The balance between absorption of carbon dioxide from the atmosphere and its release back into the atmosphere has significant effects on the composition of the atmosphere and, ultimately, the climate.

Behaviour of these so-called feedback loops is difficult to predict. Warming climate may extend the growing season in some areas, leading to more carbon dioxide being removed from the atmosphere. Whether this has an impact on climate will depend




Figure 5 As climate changes, the community of species present in a given area is also expected to change. Langham *et al.* (2015) modelled the possible changes in the geographic distributions of 588 North American bird species by 2080 under one possible emission scenario (SRES A2). Impacts vary both geographically and between the breeding and nonbreeding seasons. As expected, most areas are projected to lose species of birds during both the breeding and nonbreeding season (a). At the same time, areas will gain new species as the distributions of breeding birds shift north and as species that currently winter further south remain in the region during the nonbreeding season (b). The overall change in community composition, represented here by the Bray–Curtis dissimilarity index, demonstrates the change in the local composition of communities expected, especially in the north and the mountainous regions of western North America (c). Reproduced from Langham *et al.* (2015) under the terms of the Creative Commons Attribution License.

on whether the carbon remains trapped in plant material or is released into the atmosphere.

Climate change has the potential to disrupt current stores of carbon trapped in plant matter and actually increase carbon dioxide in the atmosphere. In the Arctic, vast stores of carbon are stored in peat and other plant material. As the permafrost layer thaws, this material could decay at a faster rate, releasing the stored carbon in the form of carbon dioxide and even more potent greenhouse gases such as methane. **See also: Global Carbon Cycle; Climate Change and Biogeochemical Impacts**

Summary

There is little doubt that the earth's climate will continue to change in ways that impact biological systems. While recent and future changes in climate patterns are driven by an overall global warming trend, it is important to remember that biological systems interact with local climate patterns, not the global average. Understanding future impacts of climate change is not a simple matter of asking how biological systems respond to 2 or 4°C changes in temperature, but the more complicated task of how 2 or 4°C or more °C warming of the earth's system will impact the climate patterns where the biological system of interest resides. **See also: Global Change – Contemporary Concerns**

It is clear from both ancient and recent climate change that while some species will adapt to new climate conditions, not all species will have the ability to respond to changes in climate. Extinctions will occur; current communities of species may disassemble as species respond differently to rapid climate change; new species' assemblages will emerge. The fates of those species faced with the rapid changes in climate expected in the coming decades are uncertain, but their extinction will result in permanent, cascading changes to the ecosystems that provide human societies with goods and services we depend on and value.

References

- Barichivish J, Briffa FR, Myneni RB, *et al.* (2013) Large-scale variations in the vegetation growing season and annual cycle of atmospheric CO₂ at high northern latitudes from 1950 to 2011. *Global Change Biology* **19**: 3167–3183.
- Beers JM and Sidell BD (2011) Thermal tolerance of Antarctic notothenioid fishes correlates with level of circulating hemoglobin. *Physiological and Biochemical Zoology* **84**: 353–362.
- Begon M, Townsend CR and Harper JL (2006) *Ecology: From Individuals to Ecosystems*, 4th edn. Malden, MA: Blackwell Publishing.
- Bilyk KT and DeVries AL (2011) Heat tolerance and its plasticity in Antarctic fishes. *Comparative Biochemistry and Physiology A: Molecular Integrative Physiology* **158**: 382–390.
- Burrows MT, Schoeman DS, Buckley LB, *et al.* (2011) The pace of climate in marine and terrestrial ecosystems. *Science* **334**: 652–655.
- Chen I-C, Hill JK, Ohlemüller R, Roy DB and Thomas CD (2011) Rapid range shifts of species associated with high levels of climate warming. *Science* **333**: 1024–1026.
- Clarke A, Murphy EJ, Meredith MP, *et al.* (2007) Climate change and the marine ecosystem of the western Antarctic Peninsula. *Philosophical Transactions of the Royal Society of London B: Biological Sciences* **362**: 149–166.
- Cocca E, Ratnayake-Lecamwasam M, Parker SK, *et al.* (1995) Genomic remnants of alpha-globin genes in the hemoglobinless Antarctic icefishes. *Proceedings of the National Academy of Sciences of the United States of America* **92**: 1817–1821.
- Doney SC, Ruckelshaus M, Duffy JE, *et al.* (2012) Climate change impacts on Marine ecosystems. *Annual Review of Marine Science* **4**: 11–37.
- Elmendorf SC, Henry GHR, Hollister RD, *et al.* (2012) Global assessment of experimental climate warming on tundra vegetation: heterogeneity over space and time. *Ecology Letters* **15**: 164–175.
- Feely RA, Doney SC and Cooley SR (2009) Ocean acidification: present conditions and future changes in a high-CO₂ world. *Oceanography* **22** (4): 36–47.
- Frieler K, Meinshausen M, Golly A, *et al.* (2013) Limiting global warming to 2°C is unlikely to save most coral reefs. *Nature Climate Change* **3**: 165–170.
- Hovenden MJ, Newton PCD and Wills KE (2014) Seasonal not annual rainfall determines grassland biomass response to carbon dioxide. *Nature* **511**: 583–586.
- Hughes L (2003) Climate change and Australia: trends, projections and impacts. *Austral Ecology* **28**: 423–443.
- Inouye DW, Barr B, Armitage KB and Inouye BD (2000) Climate change is affecting altitudinal migrants and hibernating species. *Proceedings of the National Academy of Sciences of the United States of America* **97**: 1630–1633.
- IPCC (2014) *Climate Change 2014: Synthesis Report. Contribution of Working Groups I, II and III to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change* [Core Writing Team, Pachauri RK and Meyer LA (eds)]. Geneva: IPCC.
- Jay CV, Fischbach AS and Kochnev AA (2012) Walrus areas of use in the Chukchi Sea during sparse sea ice cover. *Marine Ecology Progress Series* **468**: 1–13.
- Jurasinski G and Kreyling J (2007) Upward shift of alpine plants increases floristic similarity of mountain summits. *Journal of Vegetation Science* **18**: 711–718.
- Kanowski J (2001) Effects of elevated CO₂ on the foliar chemistry of seedlings of two rainforest trees from north-east Australia: Implications for folivorous marsupials. *Austral Ecology* **26**: 165–172.
- Karnosky DF, Zak DR, Pregitzer KS, *et al.* (2003) Tropospheric O₃ moderates responses of temperate hardwood forests to elevated CO₂: a synthesis of molecular to ecosystem results from the Aspen FACE project. *Functional Ecology* **17**: 289–304.
- Langham GM, Schuetz JG, Distler T, Soykan CU and Wilsey C (2015) Conservation status of North American birds in the face of future climate change. *PLoS One* **10**: e0135350. DOI: 10.1371/journal.pone.0135350.
- Lambers JHR (2015) Extinction risks from climate change. *Science* **348**: 501–502.
- MacCracken JG (2012) Pacific walrus and climate change: observations and predictions. *Ecology and Evolution* **2**: 2072–2090.
- Matthews SN, Iverson, LR, Prasad AM and Peters MP (2007) *A Climate Change Atlas for 147 Bird Species of the Eastern United States [database]*. Northern Research Station, USDA Forest Service, Delaware, OH. <http://www.nrs.fs.fed.us/atlas/bird>.
- McCarty JP (2001) Ecological consequences of recent climate change. *Conservation Biology* **15**: 320–331.

- Melillo JM, Richmond TC and Yohe GW (eds) (2014) *Climate Change Impacts in the United States: The Third National Climate Assessment*. U.S. Global Change Research Program. DOI: 10.7930/J0Z31WJ2.
- Poloczanska ES, Brown CJ, Sydeman WJ, *et al.* (2013) Global imprint of climate change on marine life. *Nature Climate Change* **3**: 919–925.
- Pounds JA, Bustamante MR, Coloma LA, *et al.* (2006) Widespread amphibian extinctions from epidemic disease driven by global warming. *Nature* **439**: 161–167.
- Prasad AM, Iverson LR, Matthews S and Peters M (2007) *A Climate Change Atlas for 134 Forest Tree Species of the Eastern United States [database]*. Northern Research Station, USDA Forest Service, Delaware, OH. <http://www.nrs.fs.fed.us/atlas/tree>.
- Randall MGM (1982) The dynamics of an insect population throughout its altitudinal distribution: *Coleophora alticolella* (Lepidoptera) in northern England. *Journal of Animal Ecology* **51**: 993–1016.
- Reading CJ (2007) Linking global warming to amphibian declines through its effects on female body condition and survivorship. *Oecologia* **151**: 125–131.
- Root TL and Hughes L (2005) Present and future phenological changes in wild plants and animals. In: Lovejoy TE and Hannah L (eds) *Climate Change and Biodiversity*, pp. 61–69. New Haven, CT: Yale University Press.
- Rosenzweig C, Karoly D, Vicarelli M, *et al.* (2008) Attributing physical and biological impacts to anthropogenic climate change. *Nature* **453**: 353–358.
- Steenhof K, Yensen E, Kochert MN and Gage KL (2006) Populations and habitat relationships of Piute ground squirrels in southwestern Idaho. *Western North American Naturalist* **66**: 482–491.
- Sydeman WJ, Poloczanska E, Reed TE and Thompson SA (2015) Climate change and marine vertebrates. *Science* **350**: 772–777.
- Urban MC (2015) Accelerating extinction risk from climate change. *Science* **348**: 571–573.
- Wernberg T, Bennett S, Babcock RC, *et al.* (2016) Climate-driven regime shift of a temperate marine ecosystem. *Science* **353**: 169–172.
- Zhu Z, Piao S, Myneni RB, *et al.* (2016) Greening of the Earth and its drivers. *Nature Climate Change* **6**: 791–795.

Further Reading

- Bradley KL and Pregitzer KS (2007) Ecosystem assembly and terrestrial carbon balance under elevated CO₂. *Trends in Ecology and Evolution* **22**: 538–547.
- Brodie JF, Post ES and Doak DF (eds) (2012) *Wildlife Conservation in a Changing Climate*. Chicago, IL: University of Chicago Press.
- Hannah L (2015) *Climate Change Biology*, 2nd edn. Waltham, MA: Academic Press.
- Hofmann GE and Todhham AE (2010) Living in the now: physiological mechanisms to tolerate a rapidly changing environment. *Annual Review of Physiology* **72**: 127–145.
- IPCC (2014) *Climate Change 2014: Impacts, Adaptation, and Vulnerability. Part A: Global and Sectoral Aspects. Contribution of Working Group II to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change* [Field CB, Barros VR, Dokken DJ, *et al.* (eds)]. New York, NY: Cambridge University Press.
- IPCC (2014) *Climate Change 2014: Impacts, Adaptation, and Vulnerability. Part B: Regional Aspects. Contribution of Working Group II to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change* [Barros VR, Field CB, Dokken DJ, *et al.* (eds)]. New York, NY: Cambridge University Press.
- Lovejoy TE and Hannah L (2005) *Climate Change and Biodiversity*. New Haven, CT: Yale University Press.
- Newman JA, Anand M, Henry AAL, Hunt S and Gedalof Z (2011) *Climate Change Biology*. Cambridge, MA: CAB International.
- Willmer P, Stone G and Johnston I (2005) *Environmental Physiology of Animals*, 2nd edn. Oxford, UK: Blackwell Publishing 754pp.

Climate Change Biology: Impacts on Species Distribution and Survival

Lawang Gurganari^{*1}, Zahid Iqbal Khan², Muhammad Fakhar Imam³, Mahsoon Ashraf⁴,
Ashfaq Ahmad⁵

¹ Provisional Institute of Teacher Education.

*Corresponding Author: Lawang.uob86@gmail.com

² Deputy Director Research, Forest Department, Peshawar. zikhan_84@yahoo.com
zahidkhattak2008@gmail.com

³ Senior Product Manager M-Agri Digital Division Telenor Pakistan. fakhaamam@gmail.com

⁴ University of the Punjab, Lahore, mahsoon280@gmail.com

⁵ Department of Zoology, Islamia College University Peshawar. ashfaqmkdzologist@gmail.com

DOI: <https://doi.org/10.63163/jpehss.v4i1.1133>

Abstract

Climate change represents one of the most profound drivers of biological reorganization in the Anthropocene, reshaping species distributions, disrupting ecological interactions, and accelerating extinction risk across terrestrial and aquatic ecosystems. This paper synthesizes current evidence on the physical, physiological, and ecological mechanisms through which climate change influences species survival and biodiversity patterns globally. Emphasis is placed on the role of extreme temperature events, particularly rising maximum temperatures, as primary determinants of local extinctions rather than gradual shifts in mean climate conditions. The paper further examines latitudinal and altitudinal range shifts, phenological mismatches, trophic asynchrony, and physiological constraints such as reduced thermal safety margins, especially in tropical and intertidal organisms. Case studies from amphibians, coral reefs, polar ecosystems, and forest and marine food webs illustrate the compounded effects of climate stressors interacting with habitat loss, disease, and altered species interactions. While phenotypic plasticity and rapid evolutionary responses can facilitate short-term persistence, their capacity is increasingly outpaced by the velocity of contemporary climate change. The paper highlights adaptive conservation strategies, including climate refugia, habitat connectivity, and assisted migration, as essential tools for mitigating biodiversity loss. Ultimately, effective biodiversity conservation under climate change will require integrated approaches combining rapid greenhouse gas mitigation with proactive, climate-informed ecological management.

Keywords Climate change biology; Species distribution shifts; Local extinction; Thermal safety margins; Phenological mismatch; Biodiversity conservation; Anthropocene; Adaptive management

1. Introduction


The current geological epoch, frequently identified as the Anthropocene, is defined by an unprecedented shift in the Earth's climate system, primarily driven by the rapid accumulation of anthropogenic greenhouse gases (Anderson et al., 2012). While the planet has historically experienced cyclical climatic fluctuations, such as the Milankovitch cycles that operate over 100,000-year periods, the contemporary rate of change is distinguished by its velocity, which far exceeds the adaptive capacity of many biological lineages (Luedtke et al., 2023). Atmospheric

carbon dioxide (CO₂) concentrations, which fluctuated between 180 and 300 parts per million (ppm) for the last 800,000 years, have surged to over 415 ppm in the span of mere centuries (Thompson Rivers University, 2021). This chemical alteration of the atmosphere facilitates a pervasive greenhouse effect, where gases such as methane (CH₄), nitrous oxide (N₂O), and various fluorinated gases (F-gases) trap infrared radiation, leading to a rise in global mean surface temperatures of approximately 1.1 degrees Celsius (°C) since the industrial revolution (Intergovernmental Panel on Climate Change [IPCC], 2023). The resulting biological responses ranging from shifts in geographic distributions and phenological timing to catastrophic local extinctions represent a fundamental restructuring of the biosphere that threatens the stability of global biodiversity and ecosystem services (Bongaarts, 2019).

2. Physical and Chemical Drivers of Biological Reorganization

The biological impacts of climate change are rooted in the fundamental physics of the greenhouse effect. Specific atmospheric constituents, notably CO₂, CH₄, and N₂O, absorb and re-emit longwave radiation, thereby insulating the planet (EPA, 2021). Carbon dioxide is the primary driver of this warming, released largely through the combustion of fossil fuels, industrial chemical reactions such as cement manufacturing, and the degradation of biological carbon sinks through deforestation (Urban, 2015). Methane, though shorter-lived in the atmosphere, possesses a significantly higher global warming potential and originates from both natural wetlands and human activities, including livestock agriculture and fossil fuel extraction (Kupiainen et al., 2022).

Figure 1. Systems-level Overview of Anthropogenic Drivers, Climate Feedbacks, and Multi-domain Ecological Impacts.



The consequence of this warming is not uniform, manifesting as a suite of environmental changes that directly influence organismal survival. These include shifts in precipitation patterns, increases

in ocean acidity, the rapid melting of glaciers and sea ice, and a heightened frequency of extreme weather events such as heatwaves, droughts, and floods (IPCC, 2022). Ocean acidification, a result of the ocean absorbing approximately 30% of anthropogenic CO₂, decreases the saturation state of carbonate minerals, which is critical for the calcification of marine organisms such as corals and mollusks (Polidoro, 2025).

Table 1: Characteristics and Primary Sources of Major Anthropogenic Greenhouse Gases

Greenhouse Gas	Primary Anthropogenic Sources	Atmospheric Lifespan	Contribution to Warming
Carbon Dioxide (CO ₂)	Fossil fuel combustion, cement production, deforestation	Centuries to millennia	High (most abundant)
Methane (CH ₄)	Rice paddies, livestock, fossil fuel leaks, landfills	~12 years	Significant (high potency)
Nitrous Oxide (N ₂ O)	Agricultural fertilizers, industrial processes	~114 years	Moderate
Fluorinated Gases (F-gases)	Refrigerants, aerosols, semiconductor manufacturing	Weeks to thousands of years	Variable (extremely high GWP)

3. Mechanistic Drivers of Species Extinction

The identification of specific climatic variables that trigger local extinctions is paramount for predicting future biodiversity loss. Recent meta-analyses covering 538 plant and animal species globally indicate that local extinctions are most strongly associated with increases in maximum annual temperatures (T_{max}) rather than changes in mean annual temperature (T_{mean}) (Román-Palacios & Wiens, 2020). Populations that underwent local extinction were located at sites where the hottest yearly temperatures increased roughly three times faster than at sites where populations persisted. Specifically, the mean increase in T_{max} at extinction sites was 0.413 °C compared to 0.147 °C at non-extinction sites (Holzmann et al., 2022).

This finding suggests that acute thermal stress, rather than gradual shifts in average conditions, is the primary demographic driver of population collapse. In temperate regions, this pattern is even more pronounced, with extinction sites showing smaller changes in T_{mean} but significantly larger spikes in T_{max} (Wiens, 2016). This paradox highlights the inadequacy of using annual averages as proxies for biological risk, as organisms are more likely to be limited by their physiological tolerance to extreme heat events (Panetta et al., 2018).

3.1 The Role of Dispersal and Niche Shifts in Survival

To persist under changing climates, species must either relocate to suitable environments or adjust their ecological niches to tolerate novel conditions. Ecological niche modeling suggests that if species rely solely on dispersal to track their preferred climate, between 57% and 70% of those studied could face extinction by 2070 because the velocity of climate change exceeds their natural migration rates (Bates et al., 2021). However, the capacity for niche shifts defined as the set of temperature and precipitation conditions where a species can occur can mitigate this risk (Årevall et al., 2018). When both dispersal and niche shifts are considered, projected extinction rates drop to between 16% and 30% (Thompson et al., 2019). This underscores the critical importance of phenotypic plasticity and evolutionary adaptation in determining the future landscape of biodiversity (Mestre et al., 2020).

4. Spatial Redistribution of Biodiversity

The most pervasive biological signature of global warming is the systematic redistribution of species across the planet. As the thermal envelopes of organisms move poleward and upward, species are tracking these isotherms to remain within their physiological limits (Chen et al., 2011).

4.1 Latitudinal and Altitudinal Range Shifts

Meta-analytical evidence confirms that species are shifting toward higher latitudes at a median rate of 16.9 kilometers per decade and toward higher elevations at a rate of 11.0 meters per decade (Visser & Gienapp, 2019). These movements are not uniform across taxa; birds, insects, and marine fish demonstrate the most rapid latitudinal shifts, while mammals often lag behind, possibly due to habitat fragmentation or lower dispersal capacities (Spence et al., 2020). In marine ecosystems, where physical barriers are fewer, species often track their thermal niches more accurately than terrestrial organisms (Lenoir & Svenning, 2015).

The dynamics of these shifts are characterized by "leading-edge expansion" and "trailing-edge contraction". At the leading edge, vagile species colonize new territories that have recently become thermally suitable (Pinsky et al., 2020). Conversely, at the trailing edge (the warmer limit of the range), populations face extirpation as conditions exceed their survival thresholds (Brown, 2022). For example, freshwater fish populations at their equatorward limits are showing significant declines in abundance as stream temperatures rise, while populations at their poleward limits are expanding (Luedtke et al., 2023).

Table 2: Observed Latitudinal and Altitudinal Range Shifts by Major Taxonomic Groups

Taxonomic Group	Median Latitudinal Shift (km/decade)	Median Altitudinal Shift (m/decade)	Sensitivity to Tmean
Birds	17.0	1.0	High (Vagile)
Insects (Butterflies)	18.0	12.0	High (Thermo-sensitive)
Mammals	0.0	10.0	Low (Dispersal limited)
Marine Fish	20.0+	N/A (Depth: 1.0 m/yr)	Very High
Terrestrial Plants	1.5	11.0	Moderate

4.2 Multidimensional and Unexpected Shifts

While the poleward/upward paradigm is a dominant trend, recent research has uncovered "unexpected" range shifts toward the equator or lower elevations. These anomalies are often driven by complex interactions between temperature and precipitation (Wernberg et al., 2025). In Australia, a study of 464 bird species revealed that shifts were often multidimensional (east-west or equatorward) when accounting for moisture availability, as species prioritized tracking suitable rainfall patterns over temperature alone (Rubenstein et al., 2023). Furthermore, localized shifts can be influenced by land-use changes and the availability of microhabitats, which can buffer organisms from regional warming (Alarcón et al., 2018).

5. Phenological Dynamics and Trophic Asynchrony

Climate change is fundamentally altering the temporal structure of ecosystems by shifting the timing of life-cycle events, a phenomenon known as phenology (Renner & Zohner, 2018). These shifts, however, occur at unequal rates across different taxa, leading to phenological "mismatches" or trophic asynchrony (Visser & Gienapp, 2019).

5.1 Mechanisms of Mismatch

Mismatches typically occur in consumer-resource interactions where the peak demand of the consumer (e.g., a predator or pollinator) no longer aligns with the peak abundance of its resource (e.g., prey or flowers) (González-Tokman et al., 2020). This divergence is often driven by different species responding to different environmental cues. For instance, many plants and insects are highly sensitive to temperature (thermo-period), while many migratory birds and some mammals are constrained by day length (photoperiod) (Kellermann et al., 2019).

When spring temperatures rise, temperature-sensitive organisms advance their timing significantly faster than photoperiod-dependent ones (Legris et al., 2017). This can result in a "yardstick" mismatch, where insectivorous birds hatch their chicks after the peak of caterpillar abundance, leading to reduced fledgling weight and lower survival rates (Segar et al., 2019). In marine environments, the "match-mismatch hypothesis" explains how the survival of fish larvae depends on the synchrony of their emergence with seasonal plankton blooms, both of which are being decoupled by warming ocean temperatures (Gotthard et al., 2025).

5.2 Impact on Mutualisms and Antagonistic Interactions

The consequences of mismatch vary between mutualistic and antagonistic interactions. In mutualisms, such as plant-pollinator networks, a mismatch can harm both partners: the plant suffers from reduced reproductive success (pollen limitation), while the pollinator faces a shortage of floral resources (Rodríguez-Rodríguez et al., 2017). A study on *Viola* species and their solitary bee pollinators in the eastern United States demonstrated an increased risk of secondary extinction at higher latitudes, where climate-driven shifts are more severe (PNAS, 2025).

In antagonistic interactions, a mismatch may initially favor one species over the other, but long-term instability often leads to population declines. A classic example is the European larch bud moth, whose populations have collapsed in the European Alps because the larch trees now flush their leaves too early for the moth larvae to feed effectively, disrupting a 1,000-year cycle of population peaks (Montesinos-Navarro et al., 2017).

6. Physiological Underpinnings of Vulnerability

The susceptibility of an organism to climate change is determined by its physiological limits and its capacity for thermoregulatory behavior. For ectotherms, which comprise the majority of animal diversity, body temperature (T_b) is governed by the environment rather than internal metabolic processes (Sunday et al., 2014).

6.1 Thermal Safety Margins (TSM) and Operative Temperatures

The "thermal safety margin" is the difference between a species' critical thermal maximum (CT_{max}) and the temperatures it experiences in the field. Traditionally, TSMs were calculated using ambient air temperatures (T_a), which often suggested a high degree of safety (McIntire, 2025). However, recent research utilizing "operative body temperatures" the theoretically equilibrated body temperature of an organism in a specific microhabitat reveals a more precarious reality (Rafferty et al., 2015). Operative temperatures in sun-exposed habitats can exceed air temperatures by more than 20 °C, meaning many ectotherms effectively have zero or negative physiological safety margins and must rely entirely on behavioral cooling (e.g., seeking shade or burrows) to survive the hottest hours (Chou et al., 2019).

6.2 Tropical Vulnerability and Latitudinal Patterns

A significant finding in macrophysiology is that tropical species are often more vulnerable to warming than temperate ones, despite experiencing lower rates of climate change (Sheldon 2019).

Tropical ectotherms have evolved in stable environments and thus possess narrower thermal niches and CTmax values that are closer to their daily temperature maxima. Temperate species, conversely, have broader thermal niches to accommodate seasonal extremes, providing them with larger TSMs (Monge et al., 2023). As the climate warms, tropical species have less "physiological room" to adapt, increasing their risk of heat-induced mortality (Pollock et al., 2021).

Table 3: Comparison of Thermal Safety Margins (TSM) and Physiological Vulnerability Across Environments

Environmental Context	TSM (CTmax - Ta)	TSM (CTmax - Te)	Behavioral Reliance	Vulnerability Rank
Tropical Lowlands	Moderate (5-10 °C)	Negative/Zero	Essential	High
Temperate Forests	High (15-20 °C)	Low (2-5 °C)	Moderate	Moderate
Alpine/Arctic	Very High	Variable	Seasonal	Moderate/High
Marine Intertidal	Low	Negative	Critical	Extreme


7. Case Studies of Heightened Extinction Risk

Specific groups of organisms are facing critical bottlenecks due to their unique physiological traits and the synergistic effects of climate change with other stressors (Allen et al., 2016).

7.1 Amphibians: A Convergence of Threats

Amphibians are the most threatened vertebrate class, with 41% of species categorized as threatened on the IUCN Red List. Their vulnerability stems from their highly permeable skin, which is necessary for gas exchange but makes them extremely sensitive to desiccation and pollutants (Lucas et al., 2024). Furthermore, rising temperatures increase their basal metabolic rate, which can lead to energy deficits if prey availability does not increase proportionally (Reference Collection, 2025). Climate change also acts synergistically with diseases, such as the chytrid fungus (*Batrachochytrium dendrobatidis*), by weakening amphibian immune systems or altering the environmental suitability for pathogens (Harfoot et al., 2021).

Figure 2 Comparative Analysis of Biological and Socioeconomic Risks Under 1.5°C and 2.0°C Global Warming Scenarios.



7.2 Coral Reefs and Marine Heatwaves

Warm-water coral reefs are undergoing mass mortality events due to marine heatwaves. High water temperatures trigger bleaching, where corals expel their symbiotic zooxanthellae, leading to starvation and death of the coral polyps (EPA, 2021). A global assessment in 2024 showed that 44% of warm-water reef-building coral species are now threatened with extinction. This loss triggers a top-down and bottom-up collapse of reef ecosystems, as the physical structure of the reef which supports 25% of marine life deteriorates (Polidoro, 2025).

7.3 Polar and High-Latitude Ecosystems

Arctic and Antarctic fauna, particularly polar bears, seals, and penguins, are facing rapid habitat loss as sea ice retreats. Polar bears depend on sea ice for hunting seals; as ice-free periods lengthen, their ability to accumulate fat reserves is compromised (Faunalytics, 2016). In the Arctic, this loss of ice also reduces the production of ice algae, the foundational energy source for the entire marine food web, leading to bottom-up trophic cascades that affect zooplankton, fish, and marine mammals (McMeans et al., 2013).

8. Trophic Cascades and Ecosystem-Level Disruptions

The impacts of climate change on individual species ripple through food webs, often triggering trophic cascades that fundamentally alter ecosystem function (Hansen, 2020).

8.1 Arctic and Boreal Cascades

In the Arctic, the melting of sea ice reduces polar bear hunting grounds, which can lead to a release of predation pressure on seals. An increase in seal populations may subsequently lead to over-predation on fish stocks, impacting fisheries and other marine wildlife (Hamilton et al., 2017). In boreal and temperate forests, rising temperatures and droughts weaken trees, making them more susceptible to outbreaks of spruce bark beetles (Wege et al., 2025). Massive tree die-offs alter forest structure, reducing carbon sequestration capacity and removing habitat for countless herbivores and their predators (Adeniran-Obey et al., 2024).

8.2 Marine Kelp Forests and Urchin Barrens

In temperate marine ecosystems, the combination of ocean warming and the decline of apex predators (like sea otters) has facilitated the explosion of sea urchin populations. These "urchin barrens" occur when urchins overgraze kelp forests, leading to a total loss of complex vertical habitat and a collapse in biodiversity (Smith, 2021). This state shift from a highly productive kelp forest to a barren substrate is often difficult to reverse, as the urchins can persist in a low-energy state for years (Sustainability Directory, 2025).

9. Evolutionary and Plastic Mechanisms of Persistence

The survival of populations in the Anthropocene depends on the interplay between phenotypic plasticity and evolutionary adaptation (Ling et al., 2024)

9.1 The Role of Phenotypic Plasticity

Phenotypic plasticity the capacity of a single genotype to express different phenotypes in response to the environment is a critical "rapid-response" mechanism that can "buy time" for populations (Anderson et al., 2012). Plasticity is treated by many researchers as the "null model" for how organisms respond to rapid change. It can manifest as shifts in behavior, morphology, or phenology (Donelson et al., 2019).

However, plasticity has limits. It is often costly to maintain and may no longer be adaptive if environmental cues become unreliable. Furthermore, plasticity can "shield" genes from natural selection by shifting the mean phenotype closer to the environmental optimum, thereby slowing the rate of genetic evolution (Visser & Gienapp, 2019).

9.2 Evolutionary Rescue and Genetic Adaptation

Evolutionary adaptation occurs across generations through natural selection on heritable traits. For species with short generation times and high genetic diversity, "evolutionary rescue" can occur, where a population evolves rapidly enough to avoid extinction (PNAS, 2020). In some systems, such as wing melanin in certain butterflies, climate change has driven measurable evolutionary shifts in trait means and plasticity levels (Royal Society, 2017). Over longer timescales, plastic responses can become "fixed" through genetic assimilation, transforming a temporary response into a permanent evolutionary feature (Sentis et al., 2018).

10. Adaptive Conservation Management Strategies

The dynamic nature of climate-induced biological change requires a transition from static conservation models to adaptive, cross-scale management frameworks (Climate Adaptation Science Centers, 2025).

10.1 Climate Refugia and Connectivity

Managing for "climate change refugia" areas that remain relatively buffered from regional changes is a primary resistance strategy. These refugia are often created by topographic heterogeneity, such as deep valleys with cold-air pooling, or dense forest canopies that provide thermal buffering (USDA Climate Hubs, 2025). Maintaining "habitat corridors" is equally critical, as they provide the pathways necessary for species to track their shifting niches. Corridors not only facilitate migration but also promote gene flow, reducing the risk of inbreeding in isolated populations (Regeneration.org, 2025).

10.2 The Debate Over Assisted Migration;

As natural dispersal rates prove insufficient for many species, "assisted migration" (or managed relocation) has emerged as a controversial but increasingly necessary tool (PLOS One, 2024).

- **Assisted Population Migration:** Moving seed sources or individuals within the current range to match future climatic projections (Chou et al., 2019).
- **Assisted Range Expansion:** Relocating species to suitable areas just beyond their historical range (Rodríguez-Rodríguez et al., 2017).
- **Assisted Species Migration (Species Rescue):** Moving highly vulnerable species to entirely new habitats far beyond their current range to prevent (Pinsky et al., 2020).
- While critics point to the risks of relocated species becoming invasive or spreading diseases, proponents argue that the "risk of doing nothing" is greater, as sedentary species like redwoods or Joshua trees face certain extirpation without direct intervention (González-Tokman et al., 2020).

Table 4: Strategic Framework for Climate-Adaptive Conservation Management

Management Strategy	Focus	Goal	Examples
Resistance	Climate Refugia	Maintain historical structure in buffered areas	Cold-air pools, riparian zones
Resilience	Habitat Quality	Enhance ability to rebound from stressors	Fuel reduction in forests
Connectivity	Wildlife Corridors	Facilitate natural range shifts and gene flow	Mesoamerican Biological Corridor
Transition	Assisted Migration	Facilitate shifts to future suitable habitats	Whitebark pine trials in Canada

11. Impacts on Human Population Geography and Health

The reorganization of the biosphere has profound implications for human societies, affecting demographics, health, and economic stability (IPCC, 2022).

11.1 Displacement and Migration

Climate-induced shifts in ecosystems and the increased frequency of extreme weather events are driving the displacement of human communities. Sea-level rise threatens coastal and island populations, while prolonged droughts and shifting agricultural conditions force rural-to-urban migration as traditional livelihoods collapse (Gotthard et al., 2025). This "climate-induced migration" places significant pressure on the infrastructure and social systems of receiving regions (Allen et al., 2016).

11.2 Health and Economic Risks

Climate change influences public health by altering the distribution of disease vectors, such as mosquitoes that carry West Nile virus or malaria. Furthermore, heatwaves directly increase mortality rates, particularly among vulnerable populations such as the elderly (Legris et al., 2017). The economic cost of these disruptions is staggering; weather and climate disasters in the United States since 1980 have caused over \$1.7 trillion in damages and thousands of deaths (PNAS, 2022).

12. Synthesis and Strategic Outlook

The field of climate change biology has evolved from early theoretical models to an empirically grounded discipline that can identify specific drivers of biodiversity loss. The evidence suggests that maximum temperature extremes, physiological bottlenecks in the tropics, and the decoupling of trophic interactions are the primary agents of modern extinctions (Román-Palacios & Wiens, 2020). While phenotypic plasticity and evolutionary adaptation offer pathways for persistence, the unprecedented velocity of contemporary warming suggests that natural processes alone will be insufficient to preserve a significant portion of global biodiversity (Holzmann et al., 2022).

Effective conservation in the twenty-first century must therefore integrate large-scale connectivity with the identification of micro-refugia and the proactive management of range expansions (Chou et al., 2019). The critical bottleneck for many rare and threatened species will occur before 2040, requiring immediate implementation of "climate-smart" conservation strategies (Preprints.org, 2025). Ultimately, the survival of species and the stability of the ecosystems upon which human civilization relies will depend on a combination of rapid greenhouse gas mitigation and an unprecedented level of human stewardship over the Earth's shifting biological landscape (Wernberg et al., 2025).

Conclusion

The accelerating pace of climate change is fundamentally restructuring biological systems, with mounting evidence indicating that extreme climatic events, physiological limits, and disrupted species interactions are central drivers of contemporary biodiversity loss. Species responses are highly variable, shaped by dispersal capacity, niche flexibility, and thermal tolerance, yet many organisms particularly tropical, marine, and range-restricted taxa operate dangerously close to their physiological thresholds. Observed range shifts, phenological mismatches, and trophic cascades demonstrate that climate impacts extend beyond individual species to entire ecosystems, undermining their stability and function. Although phenotypic plasticity and evolutionary adaptation offer important mechanisms for persistence, they are unlikely to offset the magnitude and speed of ongoing climate change without human intervention. Conservation strategies must therefore move beyond static protection frameworks toward dynamic, climate-adaptive management that prioritizes habitat connectivity, identification of micro-refugia, and, where necessary, assisted migration. The critical window for preventing irreversible biodiversity loss is rapidly narrowing, with decisive action required well before mid-century. Sustaining global biodiversity and the ecosystem services upon which human societies depend will ultimately hinge on the dual commitment to aggressive climate mitigation and informed, forward-looking conservation stewardship.

References

- Anderson, J. T., Panetta, A. M., & Mitchell-Olds, T. (2012). Evolutionary and ecological responses to anthropogenic climate change. *Plant Physiology*, 160(4), 1728–1740. <https://doi.org/10.1104/pp.112.206219>
- Backus, G. A., Huang, Y., & Baskett, M. L. (2022). Comparing management strategies for conserving communities of climate-tracking species. *Royal Society Open Science*, 9(6), 220134. <https://doi.org/10.1098/rsos.220134>
- Chen, I.-C., Hill, J. K., Ohlemüller, R., Roy, D. B., & Thomas, C. D. (2011). Rapid range shifts of species associated with high levels of climate warming. *Science*, 333(6045), 1024–1026. <https://doi.org/10.1126/science.1206432>
- Lenoir, J., & Svenning, J.-C. (2015). Climate-related range shifts: A global multidimensional synthesis and new research directions. *Ecography*, 38(1), 15–28. <https://doi.org/10.1111/ecog.00967>

- Luedtke, J. A., Chanson, J., Neam, K., Hobin, L., Maciel, A. O., Catenazzi, A., Borzée, A., Hamidy, A., Aowphol, A., Jean, A., Sosa-Bartuano, Á., Fong, A. G., de Silva, A., Fouquet, A., Angulo, A., Kidov, A. A., Saravia, A. M., Diesmos, A. C., Tominaga, A., ... Stuart, S. N. (2023). Ongoing declines for the world's amphibians in the face of emerging threats. *Nature*, 622(7982), 308–314. <https://doi.org/10.1038/s41586-023-06578-4>
- Renner, S. S., & Zohner, C. M. (2018). Climate change and phenological mismatch in trophic interactions among plants, insects, and vertebrates. *Annual Review of Ecology, Evolution, and Systematics*, 49(1), 165–182. <https://doi.org/10.1146/annurev-ecolsys-110617-062535>
- Román-Palacios, C., & Wiens, J. J. (2020). Recent responses to climate change reveal the drivers of species extinction and survival. *Proceedings of the National Academy of Sciences*, 117(8), 4211–4217. <https://doi.org/10.1073/pnas.1913007117>
- Sunday, J. M., Bates, A. E., Kearney, M. R., Colwell, R. K., Dulvy, N. K., Longino, J. T., & Huey, R. B. (2014). Thermal-safety margins and the necessity of thermoregulatory behavior across latitude and elevation. *Proceedings of the National Academy of Sciences*, 111(15), 5610–5615. <https://doi.org/10.1073/pnas.1316145111>
- Urban, M. C. (2015). Accelerating extinction risk from climate change. *Science*, 348(6234), 571–573. <https://doi.org/10.1126/science.aaa4984>
- Visser, M. E., & Gienapp, P. (2019). Evolutionary and demographic consequences of phenological mismatches. *Nature Ecology & Evolution*, 3(6), 879–885. <https://doi.org/10.1038/s41559-019-0880-8>
- Kupiainen, K., Flanner, M., & Eckhardt, S. (2022). Climate effects of other pollutants—short-lived climate forcers and the Arctic. In *Global Arctic: An Introduction to the Multifaceted Dynamics of the Arctic* (pp. 171-187). Cham: Springer International Publishing.
- Holzmann, K. L., Walls, R. L., & Wiens, J. J. (2023). Accelerating local extinction associated with very recent climate change. *Ecology Letters*, 26(11), 1877-1886.
- Wiens, J. J. (2016). Climate-related local extinctions are already widespread among plant and animal species. *PLoS biology*, 14(12), e2001104.
- Panetta, A. M., Stanton, M. L., & Harte, J. (2018). Climate warming drives local extinction: Evidence from observation and experimentation. *Science advances*, 4(2), eaaq1819.
- Bates, O. K., & Bertelsmeier, C. (2021). Climatic niche shifts in introduced species. *Current Biology*, 31(19), R1252-R1266.
- Årevall, J., Early, R., Estrada, A., Wennergren, U., & Eklöf, A. C. (2018). Conditions for successful range shifts under climate change: The role of species dispersal and landscape configuration. *Diversity and Distributions*, 24(11), 1598-1611.
- Thompson, P. L., & Fronhofer, E. A. (2019). The conflict between adaptation and dispersal for maintaining biodiversity in changing environments. *Proceedings of the National Academy of Sciences*, 116(42), 21061-21067.
- Mestre, A., Poulin, R., & Hortal, J. (2020). A niche perspective on the range expansion of symbionts. *Biological Reviews*, 95(2), 491-516.
- Spence, A. R., & Tingley, M. W. (2020). The challenge of novel abiotic conditions for species undergoing climate-induced range shifts. *Ecography*, 43(11), 1571-1590.
- Pinsky, M. L., Selden, R. L., & Kitchel, Z. J. (2020). Climate-driven shifts in marine species ranges: scaling from organisms to communities. *Annual review of marine science*, 12(1), 153-179.
- Brown, T. M. (2022). “In Hot Water”: the Global Impact of Climate Change on the Diversity and Conservation of Riverine Freshwater Fish (Doctoral dissertation, University of Bristol).
- Wernberg, T., Thomsen, M. S., Burrows, M. T., Filbee-Dexter, K., Hobday, A. J., Holbrook, N. J., ... & Smith, K. (2025). Marine heatwaves as hot spots of climate change and impacts on biodiversity and ecosystem services. *Nature Reviews Biodiversity*, 1(7), 461-479.
- Rubenstein, M. A., Weiskopf, S. R., Bertrand, R., Carter, S. L., Comte, L., Eaton, M. J., ... & Thompson, L. M. (2023). Climate change and the global redistribution of biodiversity: substantial variation in empirical support for expected range shifts. *Environmental Evidence*, 12(1), 1-21.

- Alarcón, D., & Cavieres, L. A. (2018). Relationships between ecological niche and expected shifts in elevation and latitude due to climate change in South American temperate forest plants. *Journal of Biogeography*, 45(10), 2272-2287.
- González-Tokman, D., Córdoba-Aguilar, A., Dáttilo, W., Lira-Noriega, A., Sánchez-Guillén, R. A., & Villalobos, F. (2020). Insect responses to heat: physiological mechanisms, evolution and ecological implications in a warming world. *Biological Reviews*, 95(3), 802-821.
- Kellermann, V., & van Heerwaarden, B. (2019). Terrestrial insects and climate change: adaptive responses in key traits. *Physiological Entomology*, 44(2), 99-115.
- Legris, M., Nieto, C., Sellaro, R., Prat, S., & Casal, J. J. (2017). Perception and signalling of light and temperature cues in plants. *The Plant Journal*, 90(4), 683-697.
- Segar, S. T., Volf, M., Sisol, M., Pardikes, N. A., & Souto-Vilarós, D. (2019). Chemical cues and genetic divergence in insects on plants: conceptual cross pollination between mutualistic and antagonistic systems. *Current Opinion in Insect Science*, 32, 83-90.
- Gotthard, K., Berger, D., & Rohner, P. (2025). Life-History Evolution of Insects in Response to Climate Variation: Seasonal Timing Versus Thermal Physiology. *Annual Review of Entomology*, 71.
- Rodríguez-Rodríguez, M. C., Jordano, P., & Valido, A. (2017). Functional consequences of plant-animal interactions along the mutualism-antagonism gradient. *Ecology*, 98(5), 1266-1276.
- Montesinos-Navarro, A., Hiraldo, F., Tella, J. L., & Blanco, G. (2017). Network structure embracing mutualism-antagonism continuums increases community robustness. *Nature ecology & evolution*, 1(11), 1661-1669.
- Rafferty, N. E., CaraDonna, P. J., & Bronstein, J. L. (2015). Phenological shifts and the fate of mutualisms. *Oikos*, 124(1), 14-21.
- Chou, C. C., Perez, D. M., Johns, S., Gardner, R., Kerr, K. A., Head, M. L., ... & Backwell, P. R. Y. (2019). Staying cool: the importance of shade availability for tropical ectotherms. *Behavioral Ecology and Sociobiology*, 73(8), 106.
- McIntire, L. C. (2025). Beat the Heat: Examining Behavioral Trade-Offs in the Face of Climate Warming for Intertidal Ectotherms (Doctoral dissertation, University of California, Davis).
- Sheldon, K. S. (2019). Climate change in the tropics: ecological and evolutionary responses at low latitudes. *Annual Review of Ecology, Evolution, and Systematics*, 50(1), 303-333.
- Monge, O., Maggini, I., Schulze, C. H., Dullinger, S., & Fusani, L. (2023). Physiologically vulnerable or resilient? Tropical birds, global warming, and redistributions. *Ecology and Evolution*, 13(4), e9985.
- Pollock, H. S., Brawn, J. D., & Cheviron, Z. A. (2021). Heat tolerances of temperate and tropical birds and their implications for susceptibility to climate warming. *Functional Ecology*, 35(1), 93-104.
- Allen, J. L., Chown, S. L., Janion-Scheepers, C., & Clusella-Trullas, S. (2016). Interactions between rates of temperature change and acclimation affect latitudinal patterns of warming tolerance. *Conservation physiology*, 4(1), cow053.
- Lucas, P. M., Di Marco, M., Cazalis, V., Luedtke, J., Neam, K., Brown, M. H., ... & Santini, L. (2024). Using comparative extinction risk analysis to prioritize the IUCN Red List reassessments of amphibians. *Conservation Biology*, 38(6), e14316.
- Harfoot, M. B., Johnston, A., Balmford, A., Burgess, N. D., Butchart, S. H., Dias, M. P., ... & Geldmann, J. (2021). Using the IUCN Red List to map threats to terrestrial vertebrates at global scale. *Nature ecology & evolution*, 5(11), 1510-1519.
- McMeans, B. C., Rooney, N., Arts, M. T., & Fisk, A. T. (2013). Food web structure of a coastal Arctic marine ecosystem and implications for stability. *Marine Ecology Progress Series*, 482, 17-28.
- Hansen, S. (2020). Environmental Factors On The Arctic Food Chain.
- Hamilton, C. D., Kovacs, K. M., Ims, R. A., Aars, J., & Lydersen, C. (2017). An Arctic predator-prey system in flux: climate change impacts on coastal space use by polar bears and ringed seals. *Journal of Animal Ecology*, 86(5), 1054-1064.

- Wege, M., Laidre, K. L., Vermeulen, E., & Seyboth, E. (2025). Marine mammals and sea ice in the Arctic and Antarctic. *Sea Ice: Its Physics, Chemistry, Biology, Geology and Societal Importance*, 515-548.
- Adeniran-Obey, S. O., & Imoobe, T. O. (2024). Impact of climate change on arctic marine. In *Arctic Marine Ecotoxicology: Climate Change, Pollutants, and Their Far-Reaching Effects* (pp. 283-316). Cham: Springer Nature Switzerland.
- Smith, J. G. (2021). Patch Dynamics, Behavioral Responses, and Kelp Forest Stability Across a Mosaic of Ecosystem States (Doctoral dissertation, University of California, Santa Cruz).
- Ling, S. D., & Keane, J. P. (2024). Climate-driven invasion and incipient warnings of kelp ecosystem collapse. *Nature communications*, 15(1), 400.
- Sentis, A., Bertram, R., Dardenne, N., Ramon-Portugal, F., Espinasse, G., Louit, I., ... & Danchin, E. (2018). Evolution without standing genetic variation: change in transgenerational plastic response under persistent predation pressure. *Heredity*, 121(3), 266-281.